

Drinking Patterns and Their Definitions (*continued*)

Low-Risk Drinking and Alcohol Use Disorder (AUD)

As defined by the National Institute on Alcohol Abuse and Alcoholism (NIAAA), for women, low-risk drinking is no more than 3 drinks on any single day and no more than 7 drinks per week. For men, it is defined as no more than 4 drinks on any single day and no more than 14 drinks per week. NIAAA research shows that only about 2 in 100 people who drink within these limits meet the criteria for AUD. Even within these limits, people can have problems if they drink too quickly or if they have other health issues.³

Binge Drinking

NIAAA defines binge drinking as a pattern of drinking that brings blood alcohol concentration to 0.08 grams per deciliter (0.08%) or higher. This typically occurs after a woman consumes 4 drinks or a man consumes 5 drinks in a 2-hour time frame.³

The Substance Abuse and Mental Health Services Administration (SAMHSA), which conducts the annual National Survey on Drug Use and Health (NSDUH), defines binge drinking as 4 or more drinks for a woman or 5 or more drinks for a man on the same occasion on at least 1 day in the past 30 days.⁴

Extreme Binge Drinking

Extreme binge drinking, also known as high-intensity drinking, refers to drinking at levels far beyond the binge threshold, resulting in high peak blood alcohol concentrations.

Though definitions vary, some studies define extreme binge drinking as 2 or more times the gender-specific binge drinking thresholds (i.e., 10 or more standard drinks for men, and 8 or more for women).⁵ Other studies use a higher threshold that may⁶ or may not⁷ be gender specific.

Heavy Drinking

SAMHSA defines heavy drinking as binge drinking on each of 5 or more days in the past 30 days.⁴

International Drink Definitions

Standard-drink definitions vary widely across countries, from 8 grams of alcohol in Iceland and the United Kingdom to 20 grams in Austria. To assess the prevalence of high-risk drinking globally, the World Health Organization uses a measure called heavy episodic drinking, defined as consuming 60 grams of alcohol or more on at least one occasion in the past 30 days. In the United States, where a standard drink equals 14 grams, that would be 4.25 standard drinks. In China, France, Ireland, and Spain, where a standard drink equals 10 grams, 6 drinks on a single occasion would constitute heavy episodic drinking.

Because of the risks of drinking, certain people should avoid alcohol completely:

- Individuals under the minimum legal drinking age of 21
- Women who are pregnant or trying to become pregnant
- People who have a medical condition that alcohol can aggravate

- Individuals taking medications that interact with alcohol
- People driving vehicles or operating machinery (or who plan to do so shortly after drinking)

References

1. What is a standard drink? National Institute on Alcohol Abuse and Alcoholism (NIAAA) Alcohol & Your Health website. <https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/what-standard-drink>. Accessed July 14, 2017.
2. *Dietary Guidelines for Americans 2015–2020*. 8th ed. Washington, DC: U.S. Department of Health and Human Services and U.S. Department of Agriculture; December 2015. <http://health.gov/dietaryguidelines/2015/guidelines>. Accessed July 14, 2017.
3. Drinking levels defined. NIAAA Alcohol & Your Health website. <https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking>. Accessed July 14, 2017.
4. Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. *Key Substance Use and Mental Health Indicators in the United States: Results from the 2015 National Survey on Drug Use and Health*. Rockville, MD: U.S. Department of Health and Human Services. <https://www.samhsa.gov/data/sites/default/files/NSDUH-FFR1-2015/NSDUH-FFR1-2015/NSDUH-FFR1-2015.pdf>. Accessed July 14, 2017.
5. Patrick ME, Crouse JM, Fairlie AM, et al. Day-to-day variations in high-intensity drinking, expectancies, and positive and negative alcohol-related consequences. *Addict Behav*. 2016;58:110-116. PMID: 26922158.
6. Patrick ME, Terry-McElrath YM. High-intensity drinking by underage young adults in the United States. *Addiction*. 2017;112(1):82-93. PMID: 27514864.
7. Johnston LD, O'Malley PM, Miech RA, et al. *Monitoring the Future National Survey Results on Drug Use, 1975–2015: 2015 Overview, Key Findings on Adolescent Drug Use*. Ann Arbor, MI: Institute for Social Research, University of Michigan; 2016. <http://monitoringthefuture.org/pubs/monographs/mf-overview2015.pdf>. Accessed July 14, 2017.